VOL. 432 JOURNAL OF CHROMATOGRAPHY NOVEMBER 18, 1988

(Biomedical Applications, Vol. 76)

CONTENTS

(Abstracts/Contents Lists published in Analytical Abstracts, ASCA, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Chromatography Abstracts, Clinical Chemistry Lookout, Current Contents/Physical, Chemical & Earth Sciences, Current Contents/Life Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, Excerpta Medica, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CNRS, Pharmaceutical Abstracts, Referativnyi Zhurnal, Science Citation Index and Trends in Biotechnology)

Analysis of lung air from patients with bronchogenic carcinoma and controls using gas chromatography–mass spectrometry
by G. Preti, J.N. Labows, J.G. Kostelc, S. Aldinger and R. Daniele (Philadelphia, PA, U.S.A.) (Received June 28th, 1988) 1

Indentification of 3-oxodicarboxylic acids in human urine
by J.S. Svendsen, L.K. Sydnes and J.E. Whist (Tromsø, Norway) (Received May 30th, 1988) 13

Determination of 17-hydroxyprogesterone in plasma by gas chromatography–mass spectrometry with high-resolution selected-ion monitoring
by K. Shimizu, T. Hara, N. Yamaga and H. Kohara (Yonago, Japan) and K. Nojima (Tokyo, Japan) (Received July 4th, 1988) 21

Rapid method for the analysis of red blood cell fatty acids by reversed-phase high-performance liquid chromatography
by G.J. Engelmann, E.L. Esmans and F.C. Alderweireldt (Antwerp, Belgium) and E. Rillaerts (Wilrijk, Belgium) (Received June 13th, 1988) 29

High-performance liquid chromatographic determination of glyoxylic acid in urine
by M. Petrarulo, S. Pellegrino, O. Bianco, M. Marangella, F. Linari and E. Mentasti (Turin, Italy) (Received June 17th, 1988) 37

Simultaneous assay for indiviual sulphated 3α- and β-hydroxysteroids in serum using high-performance liquid chromatography combined with 3α- and β-hydroxysteroid dehydrogenases immobilized on one column
by K. Takagi, K. Okumura, N. Morikawa, S. Okuyama, M.-C. Wu and O. Narita (Nagoya, Japan) (Received June 7th, 1988) 47

Investigations into lactoperoxidase-catalysed bromination of tyrosine and thyroglobulin
by W. Buchberger (Linz, Austria) (Received June 24th, 1988) 57

Determination of cysteine, glutathione and N-acetylcysteine in plasma by ion-pair reversed-phase liquid chromatography and post-column derivatization
by M. Johansson and S. Lenngren (Solna, Sweden) (Received July 7th, 1988) 65

(Continued overleaf)

Contents (continued)

Comparative chromatographic study of modifications of brush-border membrane vesicles induced by an essential fatty acid-deficient diet
by J.-M. Alessandri, R. Christon, T.S. Arfi, A. Riazi and C.L. Leger (Jouy-en-Josas, France) (Received June 3rd, 1988) 75

Affinity purification of human plasma fibronectin on immobilized gelatin
by V. Regnault, C. Rivat and J.F. Stoltz (Vandoeuvre Les Nancy, France) (Received June 28th, 1988) 93

Separation of transcobalamin II isoproteins by means of chromatofocusing
by J. Lindemans and J. van Kapel (Rotterdam, The Netherlands), F. Arwert (Amsterdam, The Netherlands) and J. Abels (Rotterdam, The Netherlands) (Received June 7th, 1988) 103

Optimization of high-performance liquid chromatography–radioimmunoassay protocols for the analyses of substances P and some of its metabolic fragments
by O.J. Igwe and L.J. Felice (St. Paul, MN, U.S.A.), V.S. Seybold (Minneapolis, MN, U.S.A.) and A.A. Larson (St. Paul, MN, U.S.A.) (Received June 14th, 1988) 113

Column-based separation of erythrocytes using aqueous polymeric two-phase systems
by D.R. Skuse and D.E. Brooks (Vancouver, Canada) (Received June 7th, 1988) 127

Free-flow electrophoretic apparatus for separation and concentration of proteins
by S.A. Shukun, A.V. Gavryushkin, V.N. Brezgunov and V.P. Zav'yalov (Moscow, U.S.S.R.) (Received July 13th, 1988) 137

Determination of clebopride in plasma by capillary gas chromatography–negative-ion chemical ionization mass spectrometry
by P.R. Robinson, M.D. Jones and J. Maddock (Merthyr Tydfill, U.K.) (Received June 6th, 1988) 153

Determination of clonidine in human plasma by cold on-column injection capillary gas chromatography–selected-ion monitoring–mass spectrometry
by R.F. Arrendale, J.T. Stewart and R.L. Tackett (Athens, GA, U.S.A.) (Received June 10th, 1988) 165

Gas chromatographic–mass spectrometric method for the quantitation of cantharidin in human serum
by J.M. Steyn and H.K.L. Hundt (Bloemfontein, Republic of South Africa) (Received June 13th, 1988) 177

Capillary gas chromatographic assay for valproic acid and its 2-desaturated metabolite in brain and plasma
by R.L.O. Semmes and D.D. Shen (Seattle, WA, U.S.A.) (Received June 28th, 1988) . 185

Determination of roxatidine in human plasma, urine and milk by capillary gas chromatography using nitrogen-selective detection
by J.L. Burrows, K.W. Jolley and D.J. Sullivan (Walton, U.K.) (Received June 16th, 1988) 199

Isolation of basic drugs from plasma using solid-phase extraction with a cyanopropyl-bonded phase
by G. Musch and D.L. Massart (Brussels, Belgium) (Received June 24th, 1988) 209

Rapid method for the determination and quantification of bromosulphophthalein and metabolites in cultured hepatocytes, culture media and bile by high-performance liquid chromatography
by G.A.E. van 't Klooster, J.H. Boot, W.C. Mennes and B.J. Blaauboer (Utrecht, The Netherlands) (Received June 30th, 1988) 223

Improved high-performance liquid chromatographic assay for the quantification of 5-bromo-2'-deoxyuridine and 5-bromouracil in plasma
by D.A. Ganes and J.G. Wagner (Ann Arbor, MI, U.S.A.) (Received May 16th, 1988) 233

Determination of bromhexine in plasma by reversed-phase liquid chromatography. Interference of lipoproteins on extraction
by M. Johansson and S. Lenngren (Solna, Sweden) (Received June 13th, 1988) 243

High-performance liquid chromatographic assay for the simultaneous monitoring of mefloquine and its acid metabolite in biological samples using protein precipitation and ion-pair extraction
by Y. Bergqvist (Falun, Sweden), U. Hellgren (Stockholm, Sweden) and F.C. Churchill (Atlanta, GA, U.S.A.) (Received June 14th, 1988) 253

Determination of atenolol in plasma by dual-column liquid chromatography and fluorimetric detection
by M. Johansson and H. Forsmo-Bruce (Solna, Sweden) (Received June 17th, 1988) . 265

Notes

Method of measurement of urinary homovanillic acid and vanillymandelic acid by gas chromatography–mass spectrometry suitable for neuroblastoma screening
by J.A. Seviour, A.C. McGill, G. Dale and A.W. Craft (Newcastle upon Tyne, U.K.) (Received June 10th, 1988) 273

Mass spectrometric analysis of endogenous diacylglycerols in normal and psoriatic skin
by A.I. Mallet, C.A. McLaughlin-Flynn and R.D.R. Camp (London, U.K.) (Received July 6th, 1988) 278

Gas chromatographic and mass spectral study of betamethasone synthetic corticosteroid metabolism
by G.M. Rodchenkov, V.P. Uralets, V.A. Semenov and V.A. Gurevich (Moscow, U.S.S.R.) (Received June 16th, 1988) 283

Gas chromatographic quantification of underivatized amphetamine in whole blood. Blood stored in plastic containers – a source of error
by A.S. Christophersen, E. Dahlin and G. Pettersen (Oslo, Norway) (Received June 13th, 1988) 290

Simple method for rapid quantification of branched-chain 2-oxo acids in physiological fluids as quinoxalinol derivatives by high-performance liquid chromatography
by W. Radeck, K. Beck and W. Staib (Düsseldorf, F.R.G.) (Received June 15th, 1988) 297

A specific reversed-phase liquid chromatographic method for analysis of steroids in Y-1 adrenal cell cultures
by B. Cheng and J. Kowal (Cleveland, OH, U.S.A.) (Received June 20th, 1988) 302

(Continued overleaf)

Contents (continued)

Simultaneous analysis of methylprednisolone hemisuccinate, cortisol and methylprednisolone by normal-phase high-performance liquid chromatography in human plasma
by A.N. Kong, R.L. Slaughter and W.J. Jusko (Buffalo, NY, U.S.A.) (Received June 28th, 1988) 308

Simultaneous determination of histidine-containing dipeptides, histamine, methylhistamine and histidine by high-performance liquid chromatography
by E. Kasziba, L. Flancbaum, J.C. Fitzpatrick, J. Schneiderman and H. Fisher (New Brunswick, NJ, U.S.A.) (Received June 22nd, 1988) 315

Estimation of tetrahydrobiopterin and other pterins in plasma by isocratic liquid chromatography with electrochemical and fluorimetric detection
by A.G. Powers, J.H. Young and B.E. Clayton (Southampton, U.K.) (Received June 17th, 1988) 321

Gas chromatographic assay for dextromoramide in human plasma
by P. Kintz, P. Mangin, A.A. Lugnier and A.J. Chaumont (Strasbourg, France) (Received June 27th, 1988) 329

Determination of maprotiline and desmethylmaprotiline in plasma and urine by gas chromatography with nitrogen–phosphorus detection
by R. Drebit, G.B. Baker and W.G. Dewhurst (Edmonton, Canada) (Received June 22nd, 1988) 334

Simplified high-performance liquid chromatographic method for determination of mizoribine in human serum
by H. Hosotsubo, S. Takahara and N. Taenaka (Osaka, Japan) (Received June 6th, 1988) 340

High-performance liquid chromatographic determination of 2′,3′-dideoxycytidine and 3′-azido-3′-deoxythymidine in plasma using a column-switching technique
by L.E. Mathes (Columbus, OH, U.S.A.), G. Muschik (Frederick, MD, U.S.A.), L. Demby and P. Polas (Columbus, OH, U.S.A.), D. Weiss Mellini and H.J. Issaq (Frederick, MD, U.S.A.) and R. Sams (Columbus, OH, U.S.A.) (Received June 30th, 1988) 346

Determination of diaziridinyl-3,6-bis(2-hydroxyethylamino)-1,4-benzoquinone in plasma by solid-phase extraction and high-performance liquid chromatography
by R.F. Betteridge, A.G. Bosanquet, P. Hunt and E.D. Gilby (Bath, U.K.) (Received June 23rd, 1988) 352

Stability of 5-aminosalicylic acid and 5-acetylaminosalicylic acid in plasma
by E. Brendel, I. Meineke, E. Stüwe and H. Osterwald (Göttingen, F.R.G.) (Received July 1st, 1988) 358

High-performance liquid chromatographic quantitation of amitriptyline and nortriptyline in dialysate from plasma or serum using on-line solid-phase extraction
by C. Svensson, G. Nyberg and E. Mårtensson (Hisings Backa, Sweden) (Received June 13th, 1988) 363

High-performance liquid chromatographic determination of chemical convulsant pentylenetetrazol in rat serum, cerebrospinal fluid and brain
by I.M. Ramzan (Pittsburgh, PA, U.S.A.) (Received June 27th, 1988) 370

High-performance liquid chromatography with fluorescence detection of dazmegrel, a specific thromboxane A_2 synthetase inhibitor
by M.V. Rebec and V.A. Skrinska (Cleveland, OH, U.S.A.) (Received July 5th, 1988) 375

Determination of oxytetracycline in fish by high-performance liquid chromatography
by H. Björklund (Åbo, Finland) (Received June 27th, 1988) 381

Assay of itraconazole in leukemic patient plasma by reversed-phase small-bore liquid chromatography
by R.P. Remmel, D. Dombrovskis and D.M. Canafax (Minneapolis, MN, U.S.A.) (Received July 4th, 1988) 388

Comparison of flow-injection analysis with high-performance liquid chromatography for the determination of etoposide in plasma
by M.A.J. van Opstal and P. Krabbenborg (Utrecht, The Netherlands), J.J.M. Holthuis (Amsterdam, The Netherlands) and W.P. van Bennekom and A. Bult (Utrecht, The Netherlands) (Received July 4th, 1988) 395

Thin-layer chromatography with flame ionization detection for the determination of tetrodotoxin in biological fluids
by J. Ikebuchi, K. Suenaga, S. Kotoku, I. Yuasa and O. Inagaki (Yonago, Japan) (Received June 21st, 1988) 401

Détermination de la céfaloglycine et de la céfroxadine dans les milieux biologiques par chromatographie sur couche mince avec détection fluorimétrique
par M.D. Blanchin et M.L. Rondot-Dudragne (Montpellier, France) (Reçu le 21 juin 1988) 407

Letter to the Editor

Simultaneous determination of floctafenine and its main metabolites, floctafenic and hydroxyfloctafenic acid, using high-performance liquid chromatography with gradient elution
by A. Tracqui, P. Kintz, P. Mangin, A.A. Lugnier and A.J. Chaumont (Strasbourg, France) (Received May 25th, 1988) 412

Related articles published in Journal of Chromatography, Vols. 449–452 415

Author Index 418

Subject Index 422

Errata 426

In articles with more than one author, the name of the author to whom correspondence should be addressed is indicated in the article heading by a 6-pointed asterisk (*)